

Comment on "Spin Polarization and Magnetic Circular Dichroism in Photoemission from the 2p Core Level of Ferromagnetic Ni"

Recently, Menchero [1] applied the four-sites cluster model [2] to the interpretation of the 2p spin-resolved x-ray photoemission spectra (SRXPS) in Ni [3]. In this Comment we show, by applying the Ni₄ cluster to the L_{2,3} magnetic circular dichroism (MCD), that the Ni ground state is not well described by this model and that it cannot provide a satisfactory description of all magnetic dichroic experiments.

The excitation of the core electron into the valence shell makes MCD more sensitive to ground-state properties than SRXPS, where the difference between the majority and minority spectrum is a result of final-state interactions between the core hole and the polarized valence shell. The calculation of the MCD spectrum, including a finite valence spin-orbit coupling (Fig. 1, dotted line), directly shows two major discrepancies between theory and experiment. First, the integrated intensities at the two spin-orbit split edges do not correspond to the experimentally observed ones. From the relations of these intensities to ground state expectation values of L_z and S_z [4,5] we find $\langle L_z \rangle / \langle S_z \rangle = 0.35$. This should be compared with the experimental value of $\langle L_z \rangle / \langle S_z \rangle = 0.19$ [5,6]. It is clear that the Ni₄ cluster overestimates the orbital magnetic moment. Second, the satellite structures, of mainly $d^8 \rightarrow \underline{p}d^9$ character, are absent.

These discrepancies are a direct result of the choice of the Ni₄ cluster. This model has a ground state consisting mainly of two holes of X_5 (t_{2g} -like) symmetry. This preference for one particular k point leads to a ground state that overestimates the orbital moment and has extremely small d^8 character, since the two holes avoid each other very effectively [2]. Choosing a two-hole ground state of X_2^2 (e_g -like, which is the second lowest state) or mixed X_2, X_5 character (by switching the energy ordering of the X_2 and X_5 band states) lowers the orbital moment to some extent, but still leads to a small d^8 character (<3%).

We were able to obtain only a ground state with a relatively small $\langle L_z \rangle / \langle S_z \rangle$ of 0.24 and a significant d^8 character (12%) by having three holes of predominantly X_2 character. Unfortunately, the good agreement for SRXPS is then lost, as Fig. 1 (solid line) shows.

In conclusion, although the Ni₄ cluster includes more information regarding the Ni band structure with respect to the Anderson impurity model [7,8], it also favors very peculiar ground states which are incompatible with a coherent picture of all dichroism experiments. Any




FIG. 1. The lower part shows the isotropic and circular dichroic L_{2,3} x-ray absorption spectra (XAS) and the upper part gives the sum and the difference of the minority (↑) and majority (↓) 2p XPS spectra, in the same geometry as in Ref. [1]. Calculations are done for the Ni₄ cluster with two X₅ (dotted) and three X₂ (solid) holes.

attempts to improve the situation by increasing the cluster size would imply formidable computational efforts.

Nicola Manini, Michel van Veenendaal,
and Massimo Altarelli
European Synchrotron Radiation Facility
B.P. 220
F-38043 Grenoble Cédex, France

Received 15 May 1997 [S0031-9007(97)04138-0]
PACS numbers: 75.25.+z, 75.30.Et, 78.20.Ls, 79.60.Bm

- [1] J. G. Menchero, Phys. Rev. Lett. **76**, 3208 (1996); Phys. Rev. B **55**, 5505 (1997).
- [2] L. M. Falicov and R. H. Victoria, Phys. Rev. B **30**, 1695 (1984); R. H. Victoria and L. M. Falicov, Phys. Rev. Lett. **55**, 1140 (1985).
- [3] A. K. See and L. E. Klebanoff, Phys. Rev. B **51**, 11002 (1995).
- [4] B. T. Thole, P. Carra, F. Sette, and G. van der Laan, Phys. Rev. Lett. **68**, 1943 (1992).
- [5] P. Carra, B. T. Thole, M. Altarelli, and X. Wang, Phys. Rev. Lett. **70**, 694 (1993).
- [6] C. T. Chen, F. Sette, Y. Ma, and S. Modesti, Phys. Rev. B **42**, 7262 (1990); J. Vogel and M. Sacchi, Phys. Rev. B **49**, 3230 (1994).
- [7] T. Jo and G. A. Sawatzky, Phys. Rev. B **43**, 8771 (1991).
- [8] G. van der Laan and B. T. Thole, J. Phys. Condens. Matter **4**, 4181 (1992).